

12-year-old boy suffers cardiac arrest due to Lyme disease

Thursday, June 01, 2017

<http://danielcameronmd.com/12-year-old-boy-suffers-cardiac-arrest-due-lyme-disease/>

by Daniel J. Cameron, MD MPH

“The patient is a 12-year-old previously healthy boy with a recent history of participation in an outdoor camp for 2–3 weeks who began to gasp for air while riding as a passenger in a car, with subsequent cyanosis and cardiac arrest, following participation in recreational outdoor activities earlier that afternoon,” [states Cunningham from the Division of Cardiology](#), Children’s National Health System, Washington, D.C.

Lyme disease triggers cardiac arrest in 12-year-old boy.

The boy received cardiopulmonary resuscitation for pulseless arrest and ventricular fibrillation. Spontaneous circulation returned “after 8 minutes of cardiopulmonary resuscitation, including defibrillation, intubation, and intravenous (IV) epinephrine and lidocaine administration,” according to Cunningham and colleagues. [1]

He required intensive therapy in the pediatric intensive care unit that included extreme hypoxemia, challenging ventilation, and persistent tachyarrhythmias including fascicular tachycardia. His treatment included high dose dopamine, norepinephrine, ceftriaxone, and procainamide. Procainamide was changed to amiodarone due to hypotension.

Serologic tests were remarkable for a strongly positive enzyme-linked immunosorbent assay (ELISA), as well as positive IgG antibodies and IgM Western blot test for Lyme disease. Intravenous ceftriaxone was continued for 21 days with normalization of his ejection fraction.

Following his participation in an outdoor camp, a 12-year-old boy suffers sudden cardiac arrest, later attributed to Lyme disease.

While the boy made a full recovery of his cardiac functioning, he did have neurologic injury which required long-term rehabilitation. Brain imaging revealed he suffered from a global cerebral edema and moderate hypoxic injury to the basal ganglia, hippocampi, and primary motor and visual cortices. The damage, Cunningham explains, left him “with altered mental status, agitation, and dysautonomia.”

The authors did not address whether the neurologic symptoms could be due to a persistent tick-borne infection yet to be identified.

Related posts:

[What might sudden cardiac death due to Lyme disease look like?](#)

[Relying on a negative Lyme disease test can prove deadly](#)

[How Lyme myocarditis might present in an adolescent patient](#)

References:

1. Cunningham MEA, Doroshov R, Olivieri L, Moak JP. Junctional ectopic tachycardia secondary to myocarditis associated with sudden cardiac arrest. HeartRhythm Case Rep. 2017;3(2):124-128.

12-year-old boy suffers cardiac arrest due to Lyme disease - <http://danielcameronmd.com/12-year-old-boy-suffers-cardiac-arrest-due-lyme-disease/>