

How to treat Babesia - a review of 38 cases at the Mayo clinic.

Sunday, October 13, 2019

<https://danielcameronmd.com/case-series-complexity-babesia/>

The authors found that nearly 50% of patients did not recall a tick bite and had a delay in diagnosis due to the broad range of symptoms. The average age was 63. None of the patients died.

Malaise was the most common symptom (84%) reported, followed by fever (71%), chills (52%) and joint pain (32%).

Sixty-eight percent of patients required hospitalization with 21% of those admitted to the intensive care unit. Indications for ICU admission included: hemolytic anemia, shock requiring vasopressor support, and acute respiratory distress syndrome.

Overall, 24% of patients were co-infected with Lyme disease; 5% with *Anaplasma* and 8% with both Lyme disease and *Anaplasma*.

“Co-infection with *Borrelia* spp. or *Anaplasma* spp. resulted in increased severity of symptoms and a higher chance of hospitalization,” [writes Fida](#).

Furthermore, 2 out of 3 patients were NOT immunocompromised.

What were the most common treatment approaches?

68% of the patients were initially treated with azithromycin and atovaquone [Mepron];

32% received clindamycin and quinine or quinidine. (*Most of the patients had to be switched from this regime to azithromycin and atovaquone due to the side effects associated with quinine.*)

Lastly, 3 severely ill patients received an exchange transfusion.

The treatment duration depended upon clinical response but ranged at least between 7 to 10 days, writes Fida. One patient, however, was treated for 6 months due to “persistently positive blood smears.” This patient was also co-infected with Lyme disease and had a previous splenectomy for leukemia.

The authors estimate that 40% of children and 20% of adults “[may experience asymptomatic infection](#),” showing no outward signs or symptoms of the disease.

Editor's note: The study looked at Babesia patients identified through the Mayo Clinic Database. Most of my Babesia cases in my practice in New York were never hospitalized.

We need additional Babesia studies that address the best treatment when other tick-borne illnesses are present. Out of those individuals hospitalized, “28.9% of patients also had Lyme disease and 10.5% had

Anaplasma co-infection,” Fida and colleagues point out.

Related Articles:

[6 cases of Babesia in early Lyme disease](#)

[Case report: Various clinical presentations of Babesia](#)

[Case series shows wide range of Babesia symptoms and presentations](#)

References:

1. Fida M, Challener D, Hamdi A, O'Horo J, Abu Saleh O. Babesiosis: A Retrospective Review of 38 Cases in the Upper Midwest. *Open Forum Infect Dis.* 2019;6(7).

Latest From the Lyme Disease Science Blog

How to treat Babesia - a review of 38 cases at the Mayo clinic. - <https://danielcameronmd.com/case-series-complexity-babesia/>