

Case series shows wide range of Babesia symptoms and presentations

Friday, February 19, 2021

<https://danielcameronmd.com/case-series-shows-wide-range-babesia-symptoms-presentations/>

This case series describes a broad range of Babesia symptoms in elderly patients, making the diagnosis particularly challenging.

Case 1: A 78-year-old white female was admitted with fever, chills, lethargy, fatigue, and marked changes in sensorium. “She had a maximum temperature of 100.6° F (38.1° C); sepsis was considered for this patient,” [writes Paparone, a Nurse Practitioner \(NP\) from the Atlantic County Health Department in Northfield, New Jersey](#). A tick-borne illness was considered, in part due to multiple tick bites, abnormal liver function tests, anemia, and thrombocytopenia. “Peripheral smear was positive for *Babesia*, and she had a *Babesia* immunoglobulin M (IgM) of 1:160 and *Anaplasma* (previously referred to as *Ehrlichia*) IgM of 1:320.” [1]

The woman’s mentation and lethargy dramatically improved when treated with a combination of doxycycline, atovaquone and zithromax.

Case 2: A 90-year-old white female was admitted for rectal bleeding with a hemoglobin of 7.6 g/dL and low platelet count of $103 \times 10^9/L$. The bleeding resolved with an octreotide infusion. The woman also had fever spikes to 100° F and a positive smear for *Babesia*. Her illness resolved with azithromycin and atovaquone.

Case 3: A 57-year-old white male was admitted with fever, malaise, and chills. His temperature had risen to 101° F during his 5-day hospital stay. *Anaplasmosis* was suspected due to his elevated liver enzymes, leukopenia, and thrombocytopenia. Intravenous doxycycline, oral clindamycin and quinine were prescribed. But he tested positive for *Babesia*. His hospital course was complicated by acute hearing deterioration. Quinine was stopped and his treatment was changed to oral azithromycin and oral atovaquone.

Case 4: An 81-year-old white male was admitted with increasing lethargy, weakness, chills, and blurred vision. *Babesia* was diagnosed on peripheral smear. *Anaplasmosis* was suspected based on anemia and thrombocytopenia.

Subsequently, serologic studies demonstrated an *Anaplasmosis* IgG of 1:256, *Babesia* IgM and IgG of 1:320. He was discharged after a 10-day combination of azithromycin and doxycycline. There was no evidence *Babesia* was treated during the hospitalization.

“At discharge on day 10, [he] was switched to clindamycin orally three times a day and quinine orally three times a day because of intolerance to azithromycin, and he completed a 14-day course of therapy,” writes Paparone.

Case 5: An 85-year-old white male was admitted with intermittent recurring fevers and chills. “He had a history of hairy cell leukemia, splenectomy, atrioventricular block (pacemaker), gouty arthritis, prostatic hypertrophy, and polymyalgia rheumatica,” writes Paparone. *Babesia* was diagnosed with 10.4% of his red blood cells infected. He was prescribed oral azithromycin and atovaquone. Doxycycline was added due to the possibility of a concurrent tick-borne infection.

He was discharged on day 8 only to be readmitted with an inability to ambulate and generalized weakness. His peripheral smear was positive for *Babesia*. “Due to the persistence of parasitemia despite adequate therapy, he was changed to clindamycin,” according to Paparone. His treatment was changed back to azithromycin and atovaquone due to gastric distress and a generalized erythematous coalescing rash. A peripheral smear for *Babesia* was negative at 5.5 weeks.

Each of the five cases presented differently:

1. Fever, chills, lethargy, fatigue, and marked changes in sensorium
2. GI bleed
3. Fever, malaise, and chills
4. Increasing lethargy, weakness, chills, and blurred vision
5. Intermittent recurring fevers and chills

Co-infections

Three of the five cases with babesia symptoms were treated for co-infections without confirmatory serologic tests. Two of three cases were treated for *Anaplasmosis* without serologic confirmation.

Treatment tolerance

Zithromax and atovaquone were well tolerated in a population of patients with babesia symptoms that included 4 elderly patients ranging from 78 to 90 years old. Quinine was stopped due to hearing loss in one subject. Clindamycin and quinine were stopped in a second subject due to gastric distress and a generalized erythematous coalescing rash.

There was no evidence any of the 5 subjects babesia symptoms required blood transfusions despite their anemia and thrombocytopenia. This suggests that prompt recognition of *Babesia* in the hospital setting might avoid the transfusions described in the literature.

Babesia was successfully treated even in their immunocompromised patient, who was treated with exchange transfusion due to persistent parasitemia. “Red blood cell exchange transfusions are recommended for cases of severe *babesiosis* in patients with parasitemia of 10% or greater, severe anemia (hemoglobin less than 10 g/dL), or pulmonary, kidney or liver impairment,” writes Paparone. “Exchange transfusions are used to rapidly decrease parasitemia, correct anemia, and help remove toxic byproducts produced by the infection.”

Authors' recommendations

“This case series illustrates the need for the NP to appreciate the variable clinical presentations of

babesiosis to facilitate prompt diagnosis, provide proper therapeutic management, and avoid the poor outcomes associated with this disease.” [1]

- It is important for the NP to understand that infected patients may not recall a tick bite and that clinical presentations may not only be variable but also nonspecific, ranging from subclinical to severe.
- The possibility of co-infection with other tick-borne illnesses (Lyme disease and anaplasmosis) must be considered.
- Furthermore, the NP needs to assume an active role in patient education to affect babesiosis awareness and prevention.

Related Articles:

[Case report: Delayed onset babesia](#)

[Geriatric babesia cases are rising and may require longer treatment](#)

[Babesia infection and tick-borne agents transmitted through blood supply](#)

References:

1. Paparone, P. and P.W. Paparone, Variable clinical presentations of babesiosis: A case series. Nurse Pract, 2017. 42(11): p. 1-7.

Latest From the Lyme Disease Science Blog

Case series shows wide range of Babesia symptoms and presentations -
<https://danielcameronmd.com/case-series-shows-wide-range-babesia-symptoms-presentations/>