

European Ixodes tick found to carry Alpha Gal protein associated with red meat allergy

Tuesday, October 17, 2017

<https://danielcameronmd.com/european-ixodes-tick-found-carry-alpha-gal-protein-associated-red-meat-allergy/>

Alpha Gal is a sugar molecule that can be spread through the bite of a Lone Star tick. Once bitten, a person's immune system begins making antibodies to α -Gal. Red meat is rife with it. So most patients become aware of the allergy or illness after they eat red meat.

[In an interview with National Geographic](#), Cosby Stone, an allergy and immunology fellow at Vanderbilt University, explained, "There's a time delay in the reaction. It [the Alpha Gal] has to first travel through your gastrointestinal tract to be released. Hours later, patients wake up with hives, shortness of breath, vomiting, and diarrhea."

European tick carries protein associated with red meat allergy

European researchers recently identified α -Gal in the gastrointestinal tract of the tick species *Ixodes (I.) ricinus*, the vector responsible for causing Lyme borreliosis (LB) in Europe. [1] Such findings suggest that tick bites from various species may lead to the development of IgE antibodies to α -Gal.

This is the first time a tick species other than the Lone Star tick has been found to carry the Alpha Gal protein. In the United States the protein has been found to trigger red meat allergies in some individuals.

According to researchers in Europe, the *I. ricinus* can transmit the protein, causing an immune response in individuals. Whether such a reaction would lead to a red meat allergy, as seen in the U.S., is unknown. The study was not designed to address "the significance of this finding in relation to red meat allergy," states Tjernberg.

The research team, led by [Tjernberg from Kalmar County Hospital in Sweden](#), looked at IgE α -Gal in 148 patients who presented with an erythema migrans (EM) rash in a separate prospective study in 2003. Serum samples were drawn at the first consultation, at 2 to 3 months and again at 6 months.

Tjernberg and colleagues found IgE anti- α -Gal reactivity was elevated. "A total of 36/148 (24%) EM patients were IgE positive to α -Gal in at least one of the three samples," writes Tjernberg and colleagues. [1]

"IgE anti- α -Gal reactivity in EM patients peaked within three months of diagnosis of EM, after which it waned indicating that recent tick exposure is of importance in α -Gal sensitization." The levels of IgE anti- α -Gal reactivity dropped over time in the general population.

The authors were able to conclude, "This indicates that recent tick exposure [to the European *I. ricinus* ticks] is of importance in sensitization to α -Gal. Furthermore, IgE anti- α -Gal was more common in men

compared with women in both blood donors and EM patients.”

References:

1. Tjernberg I, Hamsten C, Apostolovic D, van Hage M. IgE reactivity to alpha-Gal in relation to Lyme borreliosis. PLoS One. 2017;12(9):e0185723.

European Ixodes tick found to carry Alpha Gal protein associated with red meat allergy -
<https://danielcameronmd.com/european-ixodes-tick-found-carry-alpha-gal-protein-associated-red-meat-allergy/>