

Even more evidence of Lyme disease in the South

Friday, July 15, 2016

<https://danielcameronmd.com/even-more-evidence-of-lyme-disease-in-the-south/>

This expansion into new geographic territories includes regions in the South, often thought to be free of ticks carrying *Borrelia burgdorferi*, the causative agent of Lyme disease. Investigators, for instance, have identified blacklegged ticks (*Ixodes scapularis*) or western blacklegged ticks (*Ixodes pacificus*) ticks in 71 (28%) of the 254 counties in Texas. [1] Both of these species can transmit Lyme disease.

According to Rebecca Eisen, lead author of a new study, published in the *Journal of Medical Entomology*, 26 counties (10.2%) in Texas have established *I. scapularis* populations, where six or more individuals or one or more life stages were identified in a single year. Forty-five (17.7%) additional counties reported *I. scapularis* populations.

There have been studies reporting blacklegged ticks being removed from animals in Texas during the 1980's and 1990's, says Adetunji.

If there were more resources to monitor and investigate tick distribution, we might find an even greater number of counties are affected, Eisen points out. "Lack of tick records from a county -- "no records"-- does not imply that ticks are absent from that county, only that records of ticks having been collected in the county are lacking."

"The lack of routine systematic surveillance across the continental United States of ticks of public health importance hampers our ability to define their current geographic distributions and to monitor changes in their ranges and densities over time." [1]

Meanwhile, investigators led by Adetunji from the College of Veterinary Medicine & Biomedical Sciences at Texas A&M University offer additional state surveillance data. Adetunji and colleagues have identified the seroprevalence of *B. burgdorferi* antibodies (the cause of LD) in 1, 493 white-tailed deer, collected during the 2001 - 2015 hunting seasons in Texas. [2]

The findings did not surprise investigators. "A number of reports about the isolation of *B. burgdorferi* spirochetes from humans, as well as from *I. scapularis* ticks removed from animals in Texas were published during the 1980's and 1990's," reports Adetunji.

He identified six additional papers published between 2012 and 2016 which report the emergence of Lyme disease in the Texas-Mexico transboundary southern U.S. region. [2]

In the study, nearly 5% of the specimens were positive by ELISA. "Using ELISA, 4.7% of the samples were sero-reactive against *B. burgdorferi*, and these originated in two eco-regions in Texas (Edwards Plateau and South Texas Plains)," states Adetunji. Additionally, he notes that positive samples in white-tailed deer "correlate with the years in which Texas reported to CDC their highest numbers of human LD." [2]

Only 0.5% of the total samples were sero-reactive by standardized IgG western immunoblot assay. [3] This should not have come as a surprise given the poor sensitivity of the IgG western blot in actual practice, as documented in several studies conducted in the Northeastern U.S. [4]

Adetunji and colleagues voiced an additional concern with interpreting a western blot IgG in white-tailed deer. "There is a lack of standardized tests for WTD [white-tailed deer] and any information on their reactivity to specific *B. burgdorferi* antigens." [2]

An additional tick-borne illness, called southern tick-associated rash illness (STARI) is present in the South but was not addressed by the Texas research group. STARI, also known as Masters disease, was described by Dr. Edwin Masters from Missouri before his untimely death. [5] STARI is transmitted by the Lone Star tick (*Amblyomma americanum*). There is no test that would have allowed the Texas research group to study STARI.

The investigators are to be congratulated for confirming concerns that tick-borne illnesses are present in the southern U.S. "It's important for people to be aware that there may be ticks in areas where they haven't seen them previously so that they can take steps to help protect themselves and their families," states Eisen. [6]

References:

1. Eisen RJ, Eisen L, Beard CB. County-Scale Distribution of *Ixodes scapularis* and *Ixodes pacificus* (Acari: Ixodidae) in the Continental United States. *J Med Entomol*, (2016).
2. Adetunji SA, Krecek RC, Castellanos G et al. Seroprevalence of *Borrelia burgdorferi* antibodies in white-tailed deer from Texas. *Int J Parasitol Parasites Wildl*, 5(2), 168-174 (2016).
3. Feder HM, Jr., Gerber MA, Luger SW, Ryan RW. Persistence of serum antibodies to *Borrelia burgdorferi* in patients treated for Lyme disease. *Clin Infect Dis*, 15(5), 788-793 (1992).
4. Fallon BA, Keilp JG, Corbera KM et al. A randomized, placebo-controlled trial of repeated IV antibiotic therapy for Lyme encephalopathy. *Neurology*, 70(13), 992-1003 (2008).
5. Masters EJ, Grigery CN, Masters RW. STARI, or Masters disease: Lone Star tick-vectored Lyme-like illness. *Infectious disease clinics of North America*, 22(2), 361-376, viii (2008).
6. Ticks Carry Lyme Disease in Almost Half of U.S. Counties in *ScientificAmerican.com* by By Lisa Rapaport January 19, 2016 at <https://www.scientificamerican.com/article/ticks-carry-lyme-disease-in-almost-half-of-u-s-counties/>. (Last accessed 7/3/16).

Even more evidence of Lyme disease in the South - <https://danielcameronmd.com/even-more-evidence-of-lyme-disease-in-the-south/>