

Infected ticks prevalent in urban areas in the United Kingdom (UK)

Friday, October 26, 2018

<https://danielcameronmd.com/infected-ticks-prevalent-in-urban-areas-in-the-united-kingdom-uk/>

A recent article, "Assessment of the Public Health Threats Posed by Vector-Borne Disease in the United Kingdom," by Medlock and colleagues explores the growing concern Lyme disease poses to residents in the UK. [1]

“There has been an increase in the numbers of reported human cases of Lyme disease,” [writes Medlock](#) in the *International Journal of Environmental Research and Public Health*.

Between 2010 and 2017, the number of *Ixodes ricinus* ticks in the UK increased by 42%. A significant number of ticks were found to be infected with *Borrelia burgdorferi*, the bacteria causing Lyme disease. Furthermore, the prevalence of infected ticks was highest in urban areas.

“Urban woodlands and woodland edge were identified as key habitats with mean prevalence rates of *Borrelia*-infected *I. ricinus* ticks at 18%,” writes Medlock. “The highest prevalence for a single location being reported as 30% of ticks infected.”

Public health officials in the UK point out the challenge in educating people about the dangers of tick-transmitted diseases “without causing undue alarm that could hamper other advances in public health, such as the realization of the health benefits of the use of green spaces by the public,” explains Medlock.

Continuing to gather data on the distribution of *I. ricinus* ticks in urban areas is crucial, Medlock points out, as “it is possible that the management of these spaces for biodiversity could be inadvertently creating habitats for ticks, as they are designed to provide habitats for their mammal and avian hosts.”

If tick hosts, such as deer, are encouraged into urban areas, studies may need to closely examine how to manage deer populations and their movements to help minimize tick bites and the risks of Lyme disease.

Related Articles:

["Urban" ticks carry B. burgdorferi s.l. and B. miyamotoi](#)

[B. burgdorferi, the pathogen that causes Lyme disease, is widespread in New York City metro area](#)

[When should you worry about ticks in your neighborhood?](#)

[Melting pot of tick-borne pathogens found in European hedgehogs](#)

References:

1. Medlock JM, Hansford KM, Vaux AGC, Cull B, Gillingham E, Leach S. Assessment of the Public Health Threats Posed by Vector-Borne Disease in the United Kingdom (UK). *Int J Environ Res Public Health*. 2018;15(10).

Infected ticks prevalent in urban areas in the United Kingdom (UK) -

<https://danielcameronmd.com/infected-ticks-prevalent-in-urban-areas-in-the-united-kingdom-uk/>